

Sex and age-specific survival and life expectancy in a free ranging population of *Indri indri* (Gmelin, 1788).

Francesca Rolle¹, Valeria Torti¹, Daria Valente¹, Chiara De Gregorio¹, Cristina Giacomini^{1*}, Achaz von Hardenberg².

Francesca Rolle

ORCID 0000-0002-2357-1135 Twitter @FrancescaRoll9

Valeria Torti

ORCID 0000-0002-6908-1203 Twitter @valeria_torti

Daria Valente

ORCID 0000-0001-6086-5135 Twitter @DariaValente

Chiara De Gregorio

ORCID 0000-0001-7017-6181 Twitter @ChiaraDeGregor7

Cristina Giacomini

ORCID orcid.org/0000-0002-8429-7723

Achaz von Hardenberg

ORCID <https://orcid.org/0000-0002-9899-1687> Twitter: @achazhardenberg

1) Department of Life Sciences and Systems Biology, University of Turin, V. Accademia Albertina, 13 10123 Torino, Italy

2) Conservation Biology Research Group, Department of Biological Sciences, University of Chester, Parkgate Road CH1 4 BJ Chester, U.K.

*Corresponding author: cristina.giacomini@unito.it

Key words

Lemur, mortality, life span, ageing, monogamy, population ecology

Abstract

The critically endangered indri (*Indri indri*) is the largest extant lemur species and its population size is projected to decline over the next three generations due to habitat loss and climate change. Accurate information on the demographic parameters driving the population dynamics of indri is urgently needed to plan future conservation actions to secure the survival of this iconic species. We monitored and followed the life histories of 68 individually recognizable indris in 10 family groups in the Maromizaha New Protected Area (Madagascar) for 12 years. We estimated age and sex-specific survival trajectories using a Bayesian hierarchical survival model and found that the survival curves for male and female indris show a similar pattern, consistent with what found typically in primates; i.e., a high infant mortality rate which declines with age in the juvenile phase and increases again for adults. Also, life expectancies at 2 years of age (e_2), were found to be similar between the sexes (e_2 females = 7.8 years; e_2 males = 7.5 years). We suggest that the lack of strong differences in the survival patterns for male and female indris are related to the strictly monogamous system and the lack of sexual dimorphism in this species. Our study provides, for the first time, robust estimates for demographic parameters of indris and one of the very few datasets on survival trajectories available for primates.

Introduction

Predicting the population dynamics and estimating the conservation status and extinction risk of species require the long-term monitoring over multiple generations of demographic parameters in free-ranging populations (Lande 1988; Kappeler 2017). Obtaining the necessary long-term data on individual life histories may, however, result difficult when dealing with animals with a long life-span (Clutton-Brock et al. 2010). Long-term studies on the demography of non-human primates are particularly rare, not only because of the peculiar long life-span of many species but also due to the difficulties in monitoring in the wild (Pollock 1975; Bronikowski et al. 2011) and in maintaining long-term research projects

in areas often with political instabilities. As remarked by Bronikowski et al. (2011), primates represent a valuable example of a class with an average long life-expectancy but for which demographic data is scarce. The general mortality rate pattern in primates presents high infant mortality that decreases during the juvenile phase, followed by a period of age-specific increasing mortality in the mid and late life (Larson et al. 2016; Bronikowski et al. 2011; Sibly et al. 1997; Gage 1998). In addition, contrary to most other mammals, primates exhibit a longer average life expectancy relatively to their body size (Didier et al. 2016).

Mortality in primates varies across species, in accordance with their ecology and is often related to their social organization. Males and females of monogamous species tend to age at comparable rates, whereas polygynous species present differences between the sexes (Clutton-Brock et al. 2007). The mortality and rate of aging is usually considerably higher for males in species where males compete for mating opportunities (Bronikowski et al. 2011). More detailed information on juvenile mortality in primates came from studies carried out in zoos and primate centers, showing that in prosimians infants' mortality ranges between 25 and 45%, with lower values for lemuroidea (Debyser 1995). Besides, long term studies of primates in their natural habitat are needed to provide better estimate on survival and life expectancy, especially for critically endangered primates.

Indri (*Indri indri*, Gmelin 1788) is the largest lemur species and it inhabits the eastern rainforests of Madagascar, living exclusively on trees and feeding mainly on leaves and flowers (Britt et al. 2002; Borruso et al. 2021). The species occupies exclusive territories (Pollock 1975) whose defense is achieved through advertisement songs that can be heard from a 3-4 kilometers distance (Giacoma et al. 2010; Torti et al. 2013). Songs allow individuals to (i) identify the location of other groups (Giacoma et al. 2010; Gamba et al. 2016) and (ii) avoid physical conflicts and wounds, with consequent mortality (Bonadonna et al. 2017; Bonadonna et al. 2020) and (iii) communicate the reproductive potential of the individuals of the group (Torti et al. 2013; Torti et al. 2017). Indris live in family groups

composed of a monogamous and stable reproductive pair, eventually with their offspring and sub-adults up to six members (Torti et al. 2017; Bonadonna et al. 2019). Females give birth to a single offspring every 2-3 years (Quinn and Wilson 2002). Indris start singing around 1 year of age (De Gregorio et al. *in press*) and, although both sexes disperse (Gamba et al. 2013), recent work showed that males could remain in their natal group until 8 years of age, while females until 4.5 years of age (De Gregorio et al. *in press*)

Indri plays an important role in the Malagasy culture, being protected by a traditional taboo (“fady”). Through these unwritten rules the local population is culturally linked to the forest, hence, hunting or eating indri is generally avoided, as they are believed to be the ancestors of Malagasy people (Jones et al. 2008). Despite this, indri’s population size is decreasing and the rate of decline is projected to accelerate over the next three generations due to habitat loss and climate change (King et al. 2020). For these reasons, indri has been classified as Critically Endangered in the IUCN Red List of Threatened Species (King et al. 2020).

A better understanding of the demographic parameters regulating the population dynamics of indri is therefore urgently needed to support conservation activities and effective management of existing populations, and consequently guarantee the future survival of this species (Schwitzer et al. 2013). Due to the scarcity of long-term studies on the life histories of individually recognisable indris, their general life expectancy is not well known. Here we model for the first time the age-sex specific survival trajectories of indris, using a Bayesian hierarchical approach proposed by Colchero and Clark (2012) to analyse a long-term dataset on the life histories of individually recognisable indri collected over 12 years in the Maromizaha Forest, Eastern Madagascar. We specifically aim to verify if there are differences in the survival patterns of males and females in this monogamous and monomorphic species and estimate sex-specific life expectancies.

Materials and Methods

Study area

The study area corresponds to the Maromizaha New Protected Area (NAP), a primary rainforest located in the Alaotra-Mangoro Region, in the District of Moramanga, eastern-central Madagascar (18°56'49" S, 48°27'33" E). The forest covers an area of 2,150 hectares (GERP 2019) and it is part of the Ankeniheny-Zahamena Corridor (CAZ), which expands for 371,000 hectares across the eastern regions of the country. The forest extension guarantees a crucial link between the northern and southern rainforests that still persist in the country (Gamba et al. 2013). The altitude ranges between 794 and 1,224 m asl, providing the typical environment for a tropical mid-altitude evergreen rainforest with an annual average rainfall of 1779 mm (Randrianarison et al. 2015).

Maromizaha is home to a wide range of species, most of which (77%) are endemic (Randrianarison et al. 2015). Concerning the fauna, 13 lemurs, 85 birds (96% of which are endemic), 25 amphibians, 34 reptiles, and more than 800 moths species can be identified (GERP 2018; Gamba et al. 2013). Three lemur species are of particular conservation concern and classified as Critically Endangered by the IUCN red list (IUCN 2020): *Indri indri*, *Propithecus diadema*, and *Varecia variegata editorum*. It can be estimated that between 55 and 100 indri groups in total inhabit the forest (Bonadonna et al. 2017).

Indris' predators can be both aerial and terrestrial (Fitchel et al. 2002). In fact, these lemurs possess a distinct type of alarm call according to the source of potential risk (Pollock 1975; Petter & Charles-Dominique 1979; Thalmann et al. 1993; Macedonia & Stanger 1994; Maretti et al. 2010): *honks* are given when terrestrial predators are presents (e.g., fossa - *Cytoprocta ferox* and Malagasy civet, also known as striped civet or fanaloka - *fossa fossana*), while *roars* are emitted in the presence of aerial predators (e.g., Madagascar harrier-hawk - *Polyboroides radiatus*). All these predators are active in Maromizaha.

132

133 **Field data collection**

134 Following a standardised protocol (Torti et al. 2013; Bonadonna et al. 2017, 2019), data
135 were collected for 12 years, from 2009 to 2020. Monitoring is active all year round, and the
136 focal group is changed every week to ensure that all the surveyed individuals are regularly
137 under control (Bonadonna et al. 2017). All the monitored groups have been habituated to
138 human presence since 2009.

139 To limit any possible disturbance, we always made behavioural observations at a
140 distance of at least 10 meters. The animals could be identified thanks to their individual
141 external features, which are rather variable and detectable from the viewing distance: the
142 colour of the fur, the size, and the presence of any peculiar traits (Gamba et al. 2014;
143 Bonadonna et al. 2020). For each individual there is an identification card, which facilitates
144 correct identification over time (see Supplementary materials SM1). The card includes
145 name, group, sex, age, reproductive status, along with pictures of the animal and a
146 description of its main characteristics.

147 In indri males and females look the same but sing differently (De Gregorio et al. 2019a,
148 2019b). Consequently, we could not determine the sex of 13 infants that died before they
149 start singing, which occurs at around 1 year of age (De Gregorio et al. *in press*).

150 As described by Pollock (1975), indris have a short diurnal activity period, starting at dawn
151 and ending before sunset. For this reason, the observations started early in the morning
152 (6.00 a.m.) and the animals were followed and monitored in all their activities until they
153 were no longer active (Torti et al. 2013).

154

155 **Dataset**

156 This study is based on 10 individually identifiable groups of indris. The total number of
157 individuals included in the study is 68. For each individual, the following variables were

collected during the fieldwork: i) Sex (male or female); ii) Social status (reproductive or juvenile); iii) Year of birth (if known); iv) Year of death (if known); v) Presence or Absence of the individual in each year (from 2009 to 2020).

We identified four age classes, in which we grouped the individuals according to their life stage (Pereira & Altmann 1985):

1) *Infants*: individuals under 1 year of age. Infants cannot survive outside the group, can be recognized for their smaller size but it is not possible to recognize their sex.

2) *Yearlings*: individuals ranging from 1 to 2 years of age. Yearlings interact with other group members independently from their parents. Most of them start singing allowing sex identification.

3) *Juveniles*: non-reproductive individuals above 2 years of age. Juveniles typically have acquired the ability to provide for their own sustenance.

4) *Adults*: individuals of the reproductive pair in a family group.

Individuals were considered dead with certainty only if their body was found or, in the case of infants if they disappeared overnight, as they are still dependent on their mother for their survival. The disappearance of older individuals could not be assumed to be caused by death as juveniles up to the age of nine can disperse from their natal group, and we cannot trace them if they move out of the study area.

Survival analysis

Data was modelled in the open-source statistical environment R (R Core Team 2019) using the package BaSTA (Bayesian survival trajectory analysis; Colchero, Jones & Rebke 2012). BaSTA implements the Bayesian hierarchical survival model for capture-mark-recapture/resight data proposed by Colchero and Clark (2012) and it copes with datasets like ours, characterised by missing data and a large proportion of unknown ages and unknown birth and death times. The complete dataset, structured as required by the BaSTA package,

along with details can be found in Supplementary materials SM2. We compared the fit of four different possible mortality probability functions to describe the age-specific survival trajectories of indris using the *multibasta* function in BaSTA: a) the Gompertz function, assuming an exponential change in age-specific mortality (Gompertz 1825; Pletcher 1999); b) the Weibull function (Pinder III et al. 1978); c) the Logistic function (Perks 1935; Vaupel et al. 1979; Pletcher 1999) and d) the simple Exponential function (Cox & Oakes 1984). We also tested if the models would be improved by the addition of additional terms to more accurately describe the shape of the mortality curves, i.e., a Makeham structure (Pletcher 1999), which adds a constant to the mortality rate, or a bathtub structure (Siler 1979) which adds a declining Gompertz function and a constant to the basic mortality rate. Mathematical details about the four functions and their shape terms can be found in the Supporting information of Colchero, Jones and Rebke (2012). For each model, we run 4 MCMC simulations for 100,000 steps, a burn-in period of 10,001 steps, and a thinning interval of 100 steps each. The optimal combination of jump standard deviations was found by an additional initial simulation of 10,000 steps setting the *updateJumps* argument in the *multibasta* function as TRUE. We conditioned all survival models to start after 2 years of age to account for the lack of data concerning sex identification in the first years of life, while early mortality was assumed to be exponential.

As recommended by Colchero, Jones and Rebke (2012), we selected the best fitting model as the one with the lowest DIC (Deviance Information Criterion, Spiegelhalter et al. 2002) value. We compared models assuming sex differences in the model parameters (Sex-Age specific survival models) against models assuming no differences between the sexes (Age-specific survival models) and we evaluated if adding sex as a covariate improved our model fit based on their DIC values. We report the Kulbach-Leibler discrepancies (KLD; Kullback & Leibler 1951) for each pair of parameters for males and females in the best fitting Sex-Age specific survival model to measure how much sex affects each parameter in

the survival model. The KLD value ranges from 0.5 to 1 where a value of 0.5 indicates complete overlap while 1 indicates no overlap in the distribution of the parameter (McCulloch 1989). Life expectancy at age 2 (e_2) was extracted from the life table obtained with the output of the best fitting model, calculated as in Larson et al. (2016).

We estimated infant and yearling mortalities with a simple Bayesian binomial model using the *stan_glm* function in the R package *rstanarm* (Goodrich, Gabry & Brilleman 2020) as an interface to the probabilistic programming language Stan (Stan Development Team 2019). The Stan model was run using the default weakly informative normal prior for the intercept (location = 0, scale = 2.5) running 4 chains of 2000 iterations each (warmup = 1000). We used the same procedure to estimate age independent overall mortality of reproductive adults. The complete dataset and R script used for this analysis can be found on Figshare (Link in Supplementary Materials SM2).

Results

A data summary of the indri's population structure in Maromizaha Forest is reported in Table 1. Yearly group composition varied from pairs of adults without offspring to a maximum of 5 individuals (the reproductive pair, 1 or 2 offspring and/or 1-2 young joining from a different family group). Among the 68 individuals, 30 were males, 25 were females. For the remaining 13 individuals, sex is unknown because they disappeared before we could analyse their song for sex determination.

There were 22 reproductive adults in total (11 males and 11 females). In group 3MZ the adult male died in 2015 likely for the injuries reported after a physical fight with the adult male of the neighbouring group) and was substituted by another immigrant male. In group 5MZ the adult female died in 2018 after a typhoon and was as well substituted by a female joining from a different family group. Since 2009, we documented a total of 39 infants,

resulting in mean natality of 0.37 infants per group/year. Infants surviving until sex could be determined had a female-biased sex ratio of 0.67 (8/12 males/females).

In the population, 20 juveniles (13 males, 7 females) disappeared and we recorded the death of only Mika, a 4 years old female, seen wounded for unknown causes. Generally, we did not succeed in tracking juveniles after their dispersal. However, two cases have been reported of individuals moving from their native group and entering a new one in the study area:

- Case 1: Maintso born in 2010, left the natal group 1MZ in 2013, when she reached 3 years of age, she joined group 10MZ. She stayed within 10MZ without reproducing until 2014, then she moved away from the study area or died.

- Case 2: In 2014, Joeline (juvenile female, group 6MZ) dispersed when she was at least 6 years old and became the reproductive female in group 10MZ.

Variability of age at dispersal is very large, as we see that some individuals disappeared at 2 years, but others were staying in the native group at least until 8 years of age. This is true for two males in two different groups: Gibet from group 4MZ and Zafy from group 8MZ who were both born in 2012 and disappeared in 2020 at 8 years of age.

Age and sex specific survival models

We found that age specific survival is best described by a Gompertz function with a bathtub shaped curve (Table 2). Coefficient estimates for the parameters of this general model are provided in table 3.

Also, for the models including sex as a categorical variable, a Gompertz function with a bathtub shaped curve was the best fitting model, with mortality first decreasing up to 5 years of age and then increasing exponentially with age in both sexes (Table 4; Fig. 1). Parameter estimates for the best fitting age-specific survival model including sex as a covariate are provided in table 5. This model, with a DIC = 1062.54, outperforms the

previous best fitting model non assuming sex differences in survival ($DIC = 1085.58$; $DDIC = 23.04$) suggesting differences in the survival trajectories between males and females. Differences, if present, were relatively small though, with $KLDc$ values below 0.55 for all parameters with the only exception of the c parameter ($KLDc = 0.59$), indicating a marginally higher age-independent mortality in males compared to females (Table 4; Fig. 1). Estimates of life span, the maximum age individuals are known to be able to reach in the wild, was 30 years for males and 35 for females.

Life expectancies at 2 years of age were similar between the sexes (e_2 females = 7.8 years; e_2 males = 7.5 years). Infant mortality (6 out of 39 infants) was estimated to be within a 95% posterior CI going from 0.07-0.25 (median posterior estimate = 0.14). Yearling mortality (4 out of 32 individuals), ranged from 0.02-0.17 (median posterior estimate = 0.07) while overall mortality of reproductive adults (independently from age) had a median posterior estimate of 0.1 (95%CI: 0.03-0.23).

Discussion

We observed 10 habituated groups of indris from 2009 to 2020 in the Maromizaha forest, Madagascar. Group size, including always a pair of reproductive adults, varied from 2 to 5 individuals. The mean natality was 0.37 per group/year with an estimated sex ratio at one year of age of 0.67. Our age-specific survival model suggests minimal differences in the survival trajectories between males and females and similar life expectancies at 2 years of age (7.5 for males and 7.8 for females). The pattern of the survival probability curves follows what is commonly found in other primate species (Larson et al. 2016; Bronikowski et al. 2011; Sibly et al. 1997; Gage 1998). According to these studies, the survival probability is low during the first years of life. Juveniles and adults present lower mortality rates, with a constant survival probability, while mortality increases with age in the last stages of life.

Long life expectancy, along with few offspring with high mortality in a species that receive extensive parental care are common characteristics of Primates (Jones 2011). In indris, infant mortality is 0.14 and 0.07 for yearlings. This result is interesting when compared to infants' mortality of strepsirrhine primate in nature (for example, 0.30 in infant baboons, Altman 1980) and in captivity, which has been reported to range between 25 and 45%, with lower values for lemuroidea (Debiser 1995). This may indicate that indri infants' survival is among the highest in strepsirrhine, although survival estimates from captive condition may differ from the ones in wild populations. This comparative low infant mortality might be the result of Indri mothers keeping strict contact with the infant for many months. They carry the newborns until five months (Pollock 1975) and share their activities for the first eight months of life (Thalmann et al. 1993).

Further, the size of indri is also limiting success in the predation rate of the relatively small predators present in Madagascar. Another important behavioral factor that contributes to reduce mortality caused by predators, is the constant group vigilance and specialized alert behaviour, which includes vocalisations, specific for aerial and terrestrial predators (Pollock 1975; Maretti et al. 2010).

Predation is not the only cause of infant mortality. In polygamous species, social and sex competition can expose infants to other environmental threats (Bronikowski et al. 2011). We propose that Indri territoriality, monogamy and family unit social organization reduce the impact of social causes of mortality present in polygamous species, such as infanticide or aggression between females (Mitani 1990). The only other lemur species for which age-specific survival data from the wild are available is Verraux's sifaka *Propithecus verreauxi*, a species belonging to the same family (Indriidae). Verraux's sifaka mortality in the first years of life is much higher 0,48 (Richard et al. 2002) as a possible consequence of living in larger groups and of sexual competition acting in their polygamous system.

Age-specific mortality shows a similar pattern between indri and sifaka, even if lifespan appears to be shorter in *P. verreauxi* (25-30 years) in comparison to the modelling estimates we obtained in indri of 30-35 years. This long-life span is confirmed by field records from the nearby forest of Mitsinjo, where at least 1 individual is 35 years old (Rainer Dolch, personal communication). A shorter life span compared to indris is found also in baboons (*Papio cynocephalus*), which is around 27-28 years (Bronikowski et al., 2011). This species is terrestrial, has a polygamous mating system, and shows a similar mortality pattern, with high average mortality during the first and the second year of life of 0.30 (Altmann, 1980).

The general shape and pattern of the survival curves appear not to be strongly sex-related. The same sex-specific survival pattern has been documented in another monogamous primate, Azara's night monkey (*Aotus azarae*), which exhibits both biparental care and a mortality pattern comparable between males and females (Larson et al. 2016). Our own findings on mostly overlapping survival trajectories between sexes in indris can be explained by its strictly monogamous social system and lack of sexual dimorphism. In fact, survival rate is highly influenced by sex differences in species in which males and females are very dimorphic, whereas it is not when the opposite condition occurs (Bronikowski et al. 2011). Indris', with no sexual dimorphism and comparable rates of survival between the sexes, falls entirely in this frame.

An important additional factor is indri's territoriality and family unit social organization that limits social conflicts. In fact, an important difference between primates concerns the level of intra-specific competition for females that can increase males' mortality, typical in chimpanzees (*Pan troglodytes*), spider monkeys (*Ateles paniscus*), and orangutans (*Pongo pygmaeus*) in particular (Allman et al. 1998). However, male competition mostly occurs in polygynous species (among prosimians *Microcebus murinus* provides a valuable example; Fietz 1999) and indris are monogamous: the reproductive male and

female form a stable pair, with recent evidence of genetic monogamy (Bonadonna et al. 2019), despite a case of extra-pair copulation has been reported in the literature (Bonadonna et al. 2014). Monogamy can indeed provide a valuable explanation for males and females overlapping survival rates. As Bronikowski et al. (2011) argue in their comparative study, the two sexes of a monogamous species tend to age at the same rate. Conversely, in polygynous species, the ageing rates differ between the sexes.

The comparable survival pattern in males and females may also be in part related to offspring care. Species in which the mother is the one mostly responsible for offspring care are characterized by a longer life span for females, while when both parents are equally responsible for offspring care, there tends to be no significant difference in the two sexes' survival (Allman et al. 1998). To summarize with Allman's words (1998): "the sex that bears the greater burden in the care of offspring will tend to survive longer". As an example, Verraux's sifaka (*Propithecus verreauxi*) females give birth for the first time later, continue reproducing longer, and live longer than other primates (Bronikowski et al. 2011). Concerning indris, the reproductive pair defends a territory with comparable investment in territory advertisement: they participate equally to the song (equal phonation length between male and female; see Giacoma et al. 2010; De Gregorio et al. 2019a) either of advertisement or defense. Males are more active when inter-group encounters occur, and although they may therefore face a higher risk than females, territorial encounters and physical fights between neighbouring groups are extremely rare (Bonadonna et al. 2020). This is certainly a strategy to reciprocally monopolize reproduction of the male and the female of the pair, but it is also a way to supply indirect parental care.

Primate slow life histories can be an adaptive strategy when infant mortality varies greatly as a consequence of climate or social conditions (Richard et al. 2002). A work by Wright et al. (2008) conducted in the rainforest of Ranomafana reported that, for *Propithecus edwardsi*, in years when rain has been decreased during lactation months, infants of

senescent mothers died before weaning. Indris are mainly folivorous (although they can integrate their diet with fruits; Pollock 1975; Powzyk and Mowry 2003), and food is available all year long in the rainforest.

Conclusion

In conclusion, here we showed that, in accordance with what generally found in primates (Bronikowski et al. 2011) males and female indri appear to have overlapping survival and mortality trajectories. We suggest that this can be explained by the lack of sexual dimorphism and the strictly monogamous social system that characterizes this species.

Our results on indris, besides contributing to the growing availability of data on survival patterns in primates and prosimians in particular (Bronikowski et al. 2011; 2016), provide for the first time rigorous estimates of age-specific survival rates in indris. These demographic parameters represent an essential element in the development of population viability analyses, allowing a more accurate estimation of the minimum viable population size (Traill et al. 2010) necessary to guarantee the future survival of this critically endangered species.

Disclosure statement

No potential conflict of interest was reported by the authors.

Geolocation information

Madagascar, Maromizaha New Protected Area, coordinates: 18°56'49'' S; 48°27' 33''E.

Acknowledgments

This research was supported by Università degli Studi di Torino and by grants from the Parco Natura Viva – Garda Zoological Park. We thank the GERP (Groupe d'Étude et de Recherche sur les Primates de Madagascar) and Dr. Rose Marie Randrianarison for the field logistics organization and all the field guides and assistants adjuvating the data collection. Research permits have been granted from Direction des Eaux et Forêts and 'Madagascar¹⁵ National Parks' (formerly ANGAP) [2009

(N°243/09/MEF/SG/DGF/DCB.SAP/SLRSE), 2010 (N°118/10/MEF/SG/DGF/DCB.SAP/SCBSE; N°293/10/MEF/SG/DGF/DCB.SAP/SCB), 2011 (N°274/11/MEF/SG/DGF/DCB.SAP/SCB), 2012 (N°245/12/MEF/SG/DGF/DCB.SAP/SCB), 2013 (permit not required as data collection was performed by Malagasy citizens only), 2014 (N°066/14/MEF/SG/DGF/DCB.SAP/SCB), 2015 (N°180/15/MEEMF/SG/DGF/DAPT/SCBT), 2016 (N°98/16/MEEMF/SG/DGF/DAPT/SCB.Re; N°217/16/MEEMF/SG/DGF/DSAP/SCB.Re), 2017 (73/17/MEEF/SG/DGF/DSAP/SCB.RE), 2018 (91/18/MEEF/SG/DGF/DSAP/SCB.Re), 2019 (118/19/MEDD/SG/DGEF/DSAP/DGRNE; 284/19/MEDD/SG/DGEF/DSAP/DGRNE), 2019/2020 (338/19/MEDD/SG/DGEF/DSAP/DGRNE)].

References

- Allman, J., Rosin, A., Kumar, R., Hasenstaub, A. (1998). Parenting and survival in anthropoid primates: Caretakers live longer. *Proceedings of the National Academy of Sciences*, 95(12), 6866-6869. <https://doi.org/10.1073/pnas.95.12.6866>
- Altmann, J. (1980). Baboon mothers and infants. Harvard University Press, Cambridge, MA.
- Bonadonna, G., Torti, V., Randrianarison, R.M., Martinet, N., Gamba, M., Giacoma, C. (2014). Behavioral correlates of extra-pair copulation in *Indri indri*. *Primates*, 55, p. 119–123. DOI 10.1007/s10329-013-0376-0
- Bonadonna, G., Torti, V., Sorrentino, V., Randrianarison, R.M., Zaccagno, M., Gamba, M., Tan, C.L., Giacoma, C. (2017). Territory exclusivity and intergroup encounters in the indris (Mammalia: Primates: Indridae: *Indri indri*) upon methodological tuning. *The European Zoological Journal*, 84(1), p. 238-251. DOI: 10.1080/24750263.2017.1318184
- Bonadonna, G., Torti, V., De Gregorio, C., Valente, D., Randrianarison, R.M., Pozzi, L., Gamba, M., Giacoma, C. (2019). Evidence of genetic monogamy in the lemur Indri (*Indri indri*). *American Journal of Primatology*, 81(6), p. e22993. DOI: 10.1002/ajp.22993
- Bonadonna, G., Zaccagno, M., Torti, V., Valente, D., De Gregorio, C., Randrianarison, R.M., Tan, C., Gamba, M., Giacoma, C. (2020). Intra-and Intergroup Spatial Dynamics of a Pair-Living Singing Primate, *Indri indri*: A Multiannual Study of Three Indri Groups in Maromizaha Forest, Madagascar. *International Journal of Primatology*, 41, p. 224-245. <https://doi.org/10.1007/s10764-019-00127-5>
- Borruso, L., Checcucci, A., Torti, V., Correa, F., Sandri, C., Luise, D., Cavani, L., Modesto, M., Spiezio, C., Mimmo, T., Cesco, S., Di Vito, M., Bugli, F., Randrianarison, R.M., Gamba, M., Rarojoson, N.J., Avesani Zaborra, C., Mattarelli, P., Trevisi, P., Giacoma, C. (2021). I Like the Way You Eat It: Lemur (*Indri indri*) Gut Mycobiome and Geophagy. *Microbial Ecology*, 1-9. <https://doi.org/10.1007/s00248-020-01677-5>

- Britt, A., Randriamandrantonirina, N.J., Glasscock, K.D., Iambana, B.R. (2002). Diet and feeding behaviour of *Indri indri* in a low-altitude rain forest. *Folia Primatologica*, 73(5), p. 225-39. DOI: 10.1159/000067455. PMID: 12566757
- Bronikowski, A.M., Altmann, J., Brockman, D.K., Cords, M., Fedigan, L.M., Pusey, A., Stoinski, T., Morris, W.F., Strier, K.B., Alberts, S.C. (2011). Aging in the Natural World: Comparative Data Reveal Similar Mortality Patterns Across Primates. *Science*, 331 (6022), p. 1325-1328. DOI: 10.1126/science.1201571
- Bronikowski, A.M., Cords, M., Alberts, S.C., Altmann, J., Brockman, D.K., Fedigan, L.M., Pusey, A., Stoinski, T., Strier, K.B., Morris, W.F. (2016). Female and male life tables for seven wild primate species. *Scientific Data* 3, 160006. <https://doi.org/10.1038/sdata.2016.6>
- Clutton-Brock, T.H., Isvaran, K. (2007). Sex differences in ageing in natural populations of vertebrates. *Proceedings of the Royal Society B: Biological Sciences*, 274(1629), p. 3097-3104. DOI: 10.1098/rspb.2007.1138
- Clutton-Brock, T., Sheldon, B. C. (2010). The seven ages of Pan. *Science*, 327(5970), p. 1207-1208. DOI: 10.1126/science.1187796
- Colchero, F., Clark, J. S. (2012). Bayesian inference on age-specific survival for censored and truncated data. *Journal of Animal Ecology*, 81(1), p. 139-149. <https://doi.org/10.1111/j.1365-2656.2011.01898.x>
- Colchero, F., Jones, O.R., Rebke, M. (2012). BaSTA: an R package for Bayesian estimation of age-specific survival from incomplete mark-recapture/recovery data with covariates. *Methods in Ecology and Evolution*, 3(3), p. 466-470. <https://doi.org/10.1111/j.2041-210X.2012.00186.x>
- Cox, D.R., Oakes, D. (1984). *Analysis of Survival Data*. Chapman and Hall, London, UK.
- Debyser, I.W.J. (1995). Prosimian juvenile mortality in zoos and primate centers. *International Journal of Primatology*, 16(6), p. 889-907. <https://doi.org/10.1007/BF02696109>
- De Gregorio, C., Valente, D., Torti, V., Zanolli, A., Colaci Cirillo, D., Bazzano, G., Sugliano, S., Randrianarison, R.M., Giacomini, C., Gamba, M. (2019a). Song structure and sex specific features in the indris. In Proceedings of Meetings on Acoustics 177ASA (Vol. 36, No. 1, p. 010002). *Acoustical Society of America*. doi: 10.1121/2.0001021
- De Gregorio, C., Zanolli, A., Valente, D., Torti, V., Bonadonna, G., Randrianarison, R. M., Giacomini, C., Gamba, M. (2019b). Female indris determine the rhythmic structure of the song and sustain a higher cost when the chorus size increases. *Current Zoology*, 65(1), p. 89-97. <https://doi.org/10.1093/cz/zoy058>

- De Gregorio, C., Carugati, F., Estienne, V., Valente, D., Raimondi, T., Torti, V., Miaretsoa, L., Ratsimbazafy, J., Gamba, M., Giacoma, C. (*in press*). Born to sing! Song development in a singing primate. *Current Zoology*.
- Didier, E.S., MacLean, A.G., Mohan, M., Didier, P.J., Lackner, A.A., Kuroda, M. J. (2016). Contributions of nonhuman primates to research on aging. *Veterinary pathology*, 53(2), p. 277-290. <https://doi.org/10.1177/0300985815622974>
- Fietz, J. (1999). Mating system of *Microcebus murinus*. *American Journal of Primatology: Official Journal of the American Society of Primatologists*, 48(2), p. 127-133. [https://doi.org/10.1002/\(SICI\)1098-2345\(1999\)48:2<127::AID-AJP4>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1098-2345(1999)48:2<127::AID-AJP4>3.0.CO;2-4)
- Fichtel, C., Kappeler, P.M. (2002). Anti-predator behavior of group-living Malagasy primates: mixed evidence for a referential alarm call system. *Behavioral Ecology and Sociobiology*, 51, p. 262–275. <https://doi.org/10.1007/s00265-001-0436-0>
- Gamba, M., Randrianarison, R.M., Torti, V., Bounous, G., Zaborra, C.A., Jonah, R., Giacoma, C. (2013). Maromizaha: conservation and community involvement. In: Schwitzer, C., Mittermeier, R.A., Davies, N., Johnson, S., Ratsimbazafy, J., Razandramanana J., Louis Jr., E.E, Rajaobelina, S. (eds). 2013. *Lemurs of Madagascar. A Strategy for Their Conservation 2013-2016*. Bristol, UK: IUCN SSC Primate Specialist Group, Bristol Conservation and Science Foundation, and Conservation International, p. 155-156.
- Gamba, M., Torti, V., Jonah, R., Rabemananjara, Z.O., Randrianarison, R.M., Sorrentino, V., Bonadonna, G., Giacoma, C. (2013). Demografia e Dinamica di Popolazione in *Indri indri*. In: *74° Congresso Nazionale dell'Unione Zoologica Italiana*, 144(144), p. 108-108. Società dei Naturalisti e Matematici di Modena. <http://hdl.handle.net/2318/138578>
- Gamba M, Torti V, Bonadonna G, Guzzo G, Giacoma C (2014). Overlapping And Synchronization In The Song Of The Indris (*Indri indri*). In: EA Cartmill, S Roberts, H Lyn & H Cornish, ed., *The Evolution of Language: Proceedings of the 10th international conference (EvoLangX)*. p. 90-97. 14-17th April, 2014. Vienna, Austria. Singapore: World Scientific Publishing Co. ISBN: 978-981- 4603-62-1.
- Gamba, M., Torti, V., Estienne, V., Randrianarison, R. M., Valente, D., Rovara, P., Bonadonna, G., Friard, O., Giacoma, C. (2016). The indris have got rhythm! Timing and pitch variation of a primate song examined between sexes and age classes. *Frontiers in Neuroscience*, 10, p. 249. <https://doi.org/10.3389/fnins.2016.00249>

- Gage, T.B. (1998). The Comparative Demography of Primates: With Some Comments on the Evolution of Life Histories. *Annual Review of Anthropology*, 27, p. 197-221.
- GERP (2018). Rapport annuel 2018, p. 12-13. Retrieved on 17/12/2020 http://www.gerp.mg/?page_id=15
- GERP (2019). Rapport annuel 2019, p. 8-9. Retrieved on 17/12/2020 http://www.gerp.mg/?page_id=15
- Giacoma, C., Sorrentino, V., Rabarivola, C., Gamba, M. (2010). Sex differences in the song of *Indri indri*. *International Journal of Primatology*, 31(4), p. 539-551. DOI: 10.1007/s10764-010-9412-8
- Gompertz, B. (1825). On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. *Philosophical Transactions of the Royal Society of London*, 115, p. 513–583.
- Goodrich, B., Gabry, J., Ali, I., Brilleman, S. (2020). rstanarm: Bayesian applied regression modeling via Stan. R package version 2.21.1. <https://mc-stan.org/rstanarm>
- IUCN 2020. The IUCN Red List of Threatened Species. Version 2020-3. <https://www.iucnredlist.org>. Downloaded on 01/01/2021.
- Jones, J. P., Andriamarivololona, M. M., Hockley, N. (2008). The importance of taboos and social norms to conservation in Madagascar. *Conservation biology*, 22(4), 976-986. <https://doi.org/10.1111/j.1523-1739.2008.00970.x>
- Jones, J.H. (2011). Primates and the evolution of long, slow life histories. *Current Biology*, 21(18), p. R708-R717. <https://doi.org/10.1016/j.cub.2011.08.025>
- Kappeler, P.M., Cuzzo, F.P., Fichtel, C., Ganzhorn, J.U., Gursky-Doyen, S., Irwin, M.T., Ichino, S., Lawler, R., Nekaris, K.A.I, Ramanamanjato, J.B., Radespiel, U., Sauther M.L., Wright, P.C., Zimmermann, E. (2017). Long-term field studies of lemurs, lorises, and tarsiers. *Journal of Mammalogy*, 98(3), p. 661-669. DOI:10.1093/jmammal/gyx013
- King, T., Dolch, R., Randriahaingo, H.N.T., Randrianarimanana, L., Ravaloharimanitra, M. (2020). *Indri indri*. *The IUCN Red List of Threatened Species* 2020: e.T10826A115565566. <https://dx.doi.org/10.2305/IUCN.UK.2020-2.RLTS.T10826A115565566.en>. Downloaded on 30 December 2020.
- Kullback, S., Leibler, R.A. (1951). On information and suiciency. *Annals of Mathematical Statistics*, 22, p. 79–86.

- Lande, R. (1988). Genetics and demography in biological conservation. *Science*, 241(4872), 1455-1460.
- Larson, S.M., Colchero, F., Jones, O.R., Williams, L., Fernandez-Duque, E. (2016). Age and sex-specific mortality of wild and captive populations of a monogamous pair-bonded primate (*Aotus azarae*). *American Journal of Primatology*, 78(3), 315-325. <https://doi.org/10.1002/ajp.22408>
- Lemaître, J. F., Ronget, V., Tidière, M., Allainé, D., Berger, V., Cohas, A., Colchero, F., Conde, D.A., Garratt, M., Liker, A., Marais, G.A.B, Scheuerlein, A., Székely, T., Gaillard, J. M. (2020). Sex differences in adult lifespan and aging rates of mortality across wild mammals. *Proceedings of the National Academy of Sciences*, 117(15), p. 8546-8553. <https://doi.org/10.1073/pnas.1911999117>
- Macedonia J.M., Stanger K.F. (1994). Phylogeny of the Lemnidae revisited: evidence from communication signals. *Folia Primatologica*, 63, p. 1-43. <https://doi.org/10.1159/000156787>
- Maretti, G., Sorrentino, V., Finomana, A., Gamba, M., Giacomini, C. (2010). Not just a pretty song: An overview of the vocal repertoire of *Indri indri*. *Journal of Anthropological Sciences*, 88, p. 151–165.
- McCulloch, R.E. (1989). Local model influence. *Journal of the American Statistical Association*, 84, p. 473–478.
- Mitani, J. C. (1990). Demography of agile gibbons (*Hylobates agilis*). *International Journal of Primatology*, 11(5), p. 411-424.
- Nunziata, S. O., Wallenhorst, P., Barrett, M. A., Junge, R. E., Yoder, A. D., Weisrock, D. W. (2016). Population and conservation genetics in an endangered lemur, *Indri indri*, across three forest reserves in Madagascar. *International Journal of Primatology*, 37(6), 688-702. DOI 10.1007/s10764-016-9932-y
- Pereira, M. E., & Altmann, J. (1985). Development of social behavior in free-living nonhuman primates. *Nonhuman primate models for human growth and development*, p. 217-309.
- Perks, W. (1932) On Some Experiments on the Graduation of Mortality Statistics. *Journal of the Institute of Actuaries*, 63, p. 12–40. <https://www.jstor.org/stable/41137425>
- Petter J.J., Charles-Dominique P. (1979). Vocal communication in prosimians. In G.A. Doyle & R.D. Martin (eds): *The study of prosimian behaviour*, p 272-282. New York Academic Press, New York.

- Pinder III, J.E., Wiener, J.G., Smith, M.H. (1978). The Weibull distribution: a method of summarizing survivorship data. *Ecology*, 59, p. 175–179.
- Pletcher, S. (1999). Model fitting and hypothesis testing for age-specific mortality data. *Journal of Evolutionary Biology*, 12, p. 430–439.
- Pollock, J.I (1975). The social behaviour and ecology of *Indri indri*. Doctoral dissertation, University of London.
- Pollock J.I. (1975). Field Observations on *Indri indri*: A Preliminary Report. In: Tattersall I., Sussman R.W. (eds) *Lemur Biology*. Springer, Boston, MA. https://doi.org/10.1007/978-1-4684-2121-7_16
- Powzyk, J.A., Mowry, C. B. (2003). Dietary and feeding differences between sympatric *Propithecus diadema diadema* and *Indri indri*. *International Journal of Primatology*, 24(6), p. 1143-1162. <https://doi.org/10.1023/B:IJOP.0000005984.36518.94>
- Quinn, A., Wilson, D. E. (2002). *Indri indri*. *Mammalian Species*, 2002 (694), 1-5. [https://doi.org/10.1644/1545-1410\(2002\)694<0001:II>2.0.CO;2](https://doi.org/10.1644/1545-1410(2002)694<0001:II>2.0.CO;2)
- R Development Core Team (2019). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>
- Randrianarison, R.M.S., Rajaonson, A., Ralison, J.M., Rabemananjara, Z., Andrianantenaina, T.D., Rabearison, J., Ratsimbazafy, J. (2015). Local socio-economic effects of protected area conservation: The case of Maromizaha forest, Madagascar. *Madagascar Conservation & Development*, 10(2), 93-97. <http://dx.doi.org/10.4314/mcd.v10i2.8>
- Richard, A.F., Dewar, R.E., Schwartz, M., Ratsirarson, J. (2002). Life in the slow lane? Demography and life histories of male and female sifaka (*Propithecus verreauxi verreauxi*) *Journal of Zoology*, 256 (4), p. 421-436. <https://doi.org/10.1017/S0952836902000468>
- Schwitzer, C., Baker-Médard, M., Dolch, R., Golden, C., Irwin, M., Johnson, S., Patel, E., Raharivololona, B.M., Ratsimbazafy, J., Razandramanana, J., Volampeno, S. (2013). Factors Contributing to Lemur Population Decline on a National Scale, and Proposed Immediate and Longer-Term Mitigation Actions. In: Schwitzer C, Mittermeier RA, Davies N, Johnson S, Ratsimbazafy J, Razafindramanana J, Louis Jr. EE, Rajaobelina S (eds). 2013. *Lemurs of Madagascar: A Strategy for Their Conservation 2013–2016*. Bristol, UK: IUCN SSC Primate Specialist Group, Bristol Conservation and Science Foundation, and Conservation International. 185 pp.

- Sibly, R.M., Collett, D., Promislow, D.E.L., Peacock, D.J., Harvey, P.H. (1997). Mortality rates of mammals. *Journal of Zoology*, 243(1), p. 1-12. <https://doi.org/10.1111/j.1469-7998.1997.tb05751.x>
- Siler, W. (1979). A competing-risk model for animal mortality. *Ecology*, 60, p. 750–757.
- Spiegelhalter, D., Best, N., Carlin, B., Linde, A.V.D. (2002). Bayesian measures of model complexity and fit. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64, p. 583–639.
- Stan Development Team (2019). Stan Modeling Language Users Guide and Reference Manual, Version 2.25. <https://mc-stan.org>
- Thalmann, U., Geissmann, T., Simona, A., Mutschler, T. (1993). The indris of Anjanaharibe-sud, northeastern Madagascar. *International Journal of Primatology*, 14(3), p. 357-381. DOI: 0164-0291D3/0600-0357507.00
- Torti, V., Gamba, M., Rabemananjara, Z. H., Giacoma, C. (2013). The songs of the indris (Mammalia: Primates: Indridae): contextual variation in the long-distance calls of a lemur. *Italian Journal of Zoology*, 80(4), p. 596-607. DOI: 10.1080/11250003.2013.845261
- Torti, V., Bonadonna, G., De Gregorio, C., Valente, D., Randrianarison, R. M., Friard, O., Pozzi, L., Gamba, M., Giacoma, C. (2017). An intra-population analysis of the indris' song dissimilarity in the light of genetic distance. *Scientific reports*, 7(1), p. 1-12. DOI:10.1038/s41598-017-10656-9
- Traill, L. W., Brook, B. W., Frankham, R. R., & Bradshaw, C. J. (2010). Pragmatic population viability targets in a rapidly changing world. *Biological Conservation*, 143(1), 28-34.
- Vaupel, J.W, Manton, K., Stallard, E. (1979). The impact of heterogeneity in individual frailty on the dynamics of mortality. *Demography*, 16, p. 439–454.
- Wright, P., King, S.J., Baden, A., Jernvall, J. (2008). Aging in wild female lemurs: sustained fertility with increased infant mortality. In: *Primate Reproductive Aging* (Vol. 36, p. 17-28). Karger Publishers. <https://doi.org/10.1159/000137677>

Figures captions

Figure 1. Age-Sex specific survival and mortality trajectories for *Indri indri* in the Maromizaha Protected Area, Madagascar. The survival and mortality curves for males and females have been estimated following a Gompertz function with a ‘bathtub’ shape term. The coloured areas represent the 95% credibility intervals. Credibility intervals have been cut at a maximum mortality rate value of 1.0 to make the figure more readable at the lower bound. Probability density plots for the estimated parameters of the model are presented on the left.




Figure 1. Age-Sex specific survival and mortality trajectories for *Indri indri* in the Maromizaha Protected Area, Madagascar. The survival and mortality curves for males and females have been estimated following a Gompertz function with a ‘bathtub’ shape term. The coloured areas represent the 95% credibility intervals. Credibility intervals have been cut at a maximum mortality rate value of 1.0 to make the figure more readable at the lower bound. Probability density plots for the estimated parameters of the model are presented on the left.

Table 1. Summary of the data collected on 10 indri groups monitored from 2009 to 2020 in Maromizaha, Madagascar. I: infant, J: Juvenile, A: Adult; F: female; M: male; Sex of dead Infants could not be determined. *including a juvenile female immigrated from Group 1MZ.

Table 2. Ranked performances for different combinations of mortality functions and shape terms for models of age specific survival in the indri population in Maromizaha, Madagascar.

Ranked performance of the Gompertz, Weibull, Logistic and Exponential functions has been measured by the values of their Deviation Information Criterion (DIC). The lowest DIC value corresponds to the best-fitting model. Δ DIC shows the difference of DIC values between each model and the best fitting model. k is the number of estimated parameters.

Model	Shape	k	DIC	Δ DIC
Gompertz	Bathtub	6	1085.58	0.00
Weibull	Makeham	4	1106.00	20.42
Weibull	Bathtub	6	1107.03	21.45
Exponential	Simple	2	1123.86	38.28
Logistic	Bathtub	7	1128.54	42.96
Logistic	Makeham	5	1146.00	60.42
Weibull	Simple	3	1149.51	63.93
Gompertz	Simple	3	1153.71	68.13
Logistic	Simple	4	1161.76	76.18
Gompertz	Makeham	4	1162.09	76.51

Table 3. Coefficient estimates and 95% credibility intervals (CI) for the parameters of the Gompertz bathtub model (the best fitting model) for age specific survival in the indri population in Maromizaha, Madagascar. The recapture probability for 2009 is included as π .

Parameter	Estimate	Lower 95% CI	Upper 95% CI
a0	-2.12	-3.75	-1.11
a1	0.47	0.07	1.80
c	0.04	0.00	0.10
b0	-4.20	-5.82	-2.85
b1	0.03	0.00	0.09
π	0.92	0.89	0.95

Table 4. Ranked performances for different combinations of mortality functions and shape terms for models of sex and age specific survival in the indri population in Maromizaha, Madagascar.

Ranked performance of the Gompertz, Weibull, Logistic and Exponential functions has been measured by the values of their Deviation Information Criterion (DIC). The lowest DIC value corresponds to the best-fitting model. Δ DIC shows the difference of DIC values between each model and the best fitting model. k is the number of estimated parameters.

Model	Shape	DIC	Δ DIC	k
Gompertz	Bathtub	1062.54	0.00	11
Weibull	Makeham	1080.70	18.16	7
Weibull	Bathtub	1098.84	36.29	11
Logistic	Bathtub	1100.77	38.23	13
Logistic	Makeham	1102.91	40.37	9
Exponential	Simple	1113.58	51.04	3
Logistic	Simple	1126.89	64.35	7
Gompertz	Makeham	1127.40	64.85	7
Weibull	Simple	1131.31	68.77	5
Gompertz	Simple	1145.22	82.68	5

Table 5. Coefficient estimates and 95% credibility intervals (CI) for the parameters of the Gompertz bathtub model (the best fitting model) for sex and age specific survival for male (M) and female (F) indri in Maromizaha, Madagascar. KLDC (Kullback-Leibler discrepancy calibration) mean values measure the difference between sexes and ranges from 0.5 to 1. When the value is closer to 0.5 it means higher similarity. The recapture probability for 2009 is included as pi.

Parameter	Estimate		Lower 95% CI		Upper 95% CI		KLDC
	F	M	F	M	F	M	
a0	-2.15	-2.35	-3.88	-4.02	-0.81	-1.03	0.52
a1	0.71	0.80	0.08	0.08	2.14	2.22	0.51
c	0.04	0.06	0.00	0.00	0.10	0.13	0.59
b0	-4.12	-3.97	-5.74	-5.62	-2.77	-2.60	0.51
b1	0.04	0.05	0.00	0.00	0.12	0.16	0.55
pi	0.92		0.89		0.95		

906 SM1. Example of an identification card with name, group, sex, age, reproductive status,
907 pictures and description of Jery, the adult reproductive male of Group 1.

908
909

Name: Jery - Group: 1MZ - Sex: Male - Age: Adult - Class: Reproductive



Reproductive male, adult. Sings in duets with *Bevolo*. Easily identifiable by the very dark colour of its fur, both in the dorsal and the ventral part of the body. The white dorsal triangle is very narrow and mottled with gray. The white parts of the fur on the forelimbs and hind limbs are mottled with smoke gray. The white triangle on the head presents two bits, not very long, that descend on the eyes, and a dark linear spot that cuts through the triangle in half, starting from the dorsal portion. The face is very dark, with a lighter and slightly perceptible halo around the eyes

910
911
912
913
914
915
916

SM 2. Data availability

In order to satisfy the requirements of the BaSTA package (Colchero, Jones and Rebke 2012), the data was structured as a matrix, where each row corresponds to an individual (ID - first column). Then, the matrix includes birth and death years (second and third column, if available) and the individual recapture history (from the fourth column onwards). The recapture history is expressed through a matrix that assigns 1 for the years when the individual was present and 0 if not observed in that year. Unknown birth and death years were coded with 0. The R Script used for this analysis and the complete dataset can be downloaded as a csv file (Data_Indri_Population.csv) from

<https://figshare.com/s/8d813f028a59f4b3db9b>

References

Colchero, F., Jones, O.R., Rebke, M. (2012). BaSTA: an R package for Bayesian estimation of age-specific survival from incomplete mark-recapture/recovery data with covariates. *Methods in Ecology and Evolution*, 3(3), p. 466-470. <https://doi.org/10.1111/j.2041-210X.2012.00186.x>